

**COP1 CONTRIBUTES TO UVB-INDUCED SIGNALING
IN HUMAN KERATINOCYTES**

Ph.D. Dissertation

Ágnes Kinyó

Supervisor:

Prof. Dr. Lajos Kemény

**Department of Dermatology and Allergology
University of Szeged, Szeged, Hungary**

2011

1. INTRODUCTION

1.1 COP1, the Constitutive Photomorphogenic Protein 1

1.1.1. Definition

UVB light, one of the most important physical carcinogens in the environment, has been shown to trigger a broad range of changes in gene expression in plants and in non-plant multicellular organisms. Many components of the light signaling cascades have been revealed over the years, mostly by genetic approaches. Among them, COP1 (constitutive photomorphogenic protein 1) was one of the first cloned and is one of the most extensively studied. COP1 is a RING-finger protein; it was first described in *Arabidopsis thaliana*.

1.1.2. COP1 in plants

It has been established that *Arabidopsis thaliana* COP1 (AtCOP1) functions as an essential negative regulator of light-mediated plant development, as an E3 ubiquitin ligase, is essential for the proteasome-dependant degradation of a number of bZIP-type transcription factors, such as HY5, HYH, LAF1 and HFR1. Genome-wide microarray analysis has demonstrated that AtCOP1 regulates most of the light-responsive genes under various light conditions. AtCOP1 functions as a crucial developmental switch through targeting key transcription factors for degradation, thereby controlling light-responsive gene expression and photomorphogenic development. In photosynthesizing cells, the subcellular distribution of AtCOP1 is adjusted according to the light conditions.

1.1.3. COP1 in non-plant organisms

Although, COP1 also exists in non-plant multicellular organisms, its function is less understood among vertebrates, COP1 is well conserved among fish, amphibians, birds and mammals, it is apparent that, unlike plants, animals do not undergo photomorphogenesis. Several initial studies have implied that animal COP1 may have adopted roles in tumorigenesis and stress response, although its biochemical activities and some of the signaling elements it interacts with appear to be conserved between plants and animals.

1.1.4. Structural and functional analogy between AtCOP1 and mammalian COP1

Mammalian COP1 (MmCOP1), as human COP1 (huCOP1), is located at the chromosome 1 and exhibits a high degree of sequence conservation and domain organization with AtCOP1, suggesting the possibility of functional conservation. Both MmCOP1 and AtCOP1 contain three conserved structural domains: a RING finger at the amino terminus, a coiled-coil domain in the middle, and a carboxyl-terminal WD40 repeat domain. Earlier studies suggested that COP1 acts primarily as a homodimer, and probably dimerizes through the coiled-coil domain. The coiled-coil region and the seven WD40 motifs region comprise potential interaction domains for other protein partners.

As compared with AtCOP1, MmCOP1 has an N-terminal extension that may be responsible for targeting COP1 to the nuclear envelope (NE) part of the nucleus. COP1 is predominantly localized in the nucleus, but a small amount may also be present in the cytosol. COP1 shuttles between the nucleus and the cytoplasm and forms subnuclear speckles in both plants and mammals.

1.2. Function of mammalian COP1

MmCOP1, like AtCOP1, is involved in ubiquitination and is itself a substrate of its own ubiquitination activity. COP1 ubiquitination substrates have been identified in mammals: c-Jun and p53. COP1-mediated p53 degradation is an important regulatory mechanism for the p53 function in the cell: depletion of COP1 leads to p53 accumulation and cell-cycle arrest, while the overexpression of COP1 inhibits p53-dependent apoptosis. Overexpression of huCOP1 in mammalian cells downregulates c-Jun-dependent transcription and expression of the AP-1 target genes.

1.3. p53 regulation in keratinocytes and in the epidermis

p53, a nuclear phosphoprotein first identified by Lane and Crawford in 1979, activates the transcription of many genes involved in cell differentiation, proliferation, induction of DNA repair pathways and cell death. The loss of p53 function increases the risk of development for certain types of cancers. As regards UV-induced skin tumors, p53 is considered to play an important role in their pathogenesis.

In normal unstressed cells, p53 is a very unstable protein with a half-life ranging from 5 to 30 min. Its short half-life is linked to polyubiquitination and proteasome-dependent degradation, while the protein is regulated through posttranslational modifications including phosphorylation and acetylation. The p53 protein level is increased and stabilized in both murine and human keratinocytes when treated with UVB light. p53 accumulates intracellularly up to readily detectable levels, arresting cells in the G1 phase, until cellular damage is fully repaired or the cell-death pathway is initiated.

In normal human skin, p53 is generally not detected in the epidermis, which has been attributed to its relatively short half-life. p53 is the key UV-responsive gene in skin, whose mutation is thought to initiate carcinogenesis, but its role in the control of keratinocyte proliferation and differentiation is less clear.

1.4. E3 ubiquitin ligases and p53 regulation

p53 expression has been shown to be controlled by a plethora of upstream regulatory proteins, the main antagonists of p53 are Mdm2, Pirh2 and huCOP1 E3 ubiquitin ligases. These proteins directly promote p53 ubiquitination and its proteasome-dependent degradation. Mdm2 interacts physically with and ubiquitinates p53, leading to its degradation in the cytosolic 26S proteasome, but it has also been demonstrated that p53 degradation may occur in either the cytosolic or the nuclear proteasomes and does not require nuclear export. In addition to Mdm2, Pirh2 and COP1 have been shown to impart specificity toward p53. The nuclear monoubiquitination of p53 by Mdm2 leads to the accumulation of p53. Following monoubiquitination, p53 is polyubiquitinated by Mdm2, Pirh2, COP1 or a complex of these proteins, and degraded by cytosolic 26S proteasomes. It has been demonstrated that COP1 promotes p53 turnover independently of Mdm2 or Pirh2 in U2OS osteosarcoma cells.

1.5. Kinetics of p53 expression during keratinocyte differentiation

Keratinocytes undergo a complex developmental program as they progress from the basal to the spinous, granular and finally the cornified layer of the epidermis. p53 is expressed at low levels in UV-unexposed skin and is activated during the exit from the proliferative state, resulting in inhibition of cell division and stimulation of differentiation. Later, as the keratinocytes progress in the differentiation associated the cell death program, they become insensitive to p53-induced apoptosis. UV light induces p53 in the proliferative compartment of the basal layer cells, resulting in cell cycle arrest and apoptosis, which protects from cancer induction. When p53 is inhibited in the basal layer by a negative regulator protein, such as Mdm2, this leads to decreased differentiation and increased proliferation.

1.6. Effects of genotoxic stress on E3 ubiquitin ligases

In wild-type p53 expressing cells, E3 ubiquitin ligases, such as Mdm2, Pirh2 and COP1, are induced following genotoxic stress, caused, for example, by ionizing radiation (IR) or UV irradiation. Mdm2 expression increases in a p53-dependent manner following IR, and its regulation after UVB irradiation in human keratinocytes is also well elucidated. Pirh2 expression is not enhanced following IR or UV treatment in wild-type p53-expressing cell lines, but it has not been investigated in human keratinocytes. COP1 protein expression increases following treatment with IR in wild-type p53 expressing cells, UVC results in biphasic mRNA changes and an elevated COP1 protein level in U2OS osteosarcoma cells and HeLa cells.

1.7. COP1 in human tumors

As COP1 is a critical negative regulator of p53 in normal and cancer cell lines, its role has been investigated in detail in human cancers. The overexpression of COP1 was detected predominantly, but not exclusively, in wild-type p53-containing cancers, such as ovarian tumors, breast adenocarcinomas and hepatocellular carcinomas, indicating that one of the major roles of COP1 is to repress p53-dependent tumor suppression. Finally, COP1 may function as an oncogene and promote tumorigenesis, especially in tumors in which p53 is not mutated. Investigation of COP1 and the p53-associated functional network in hepatocellular carcinomas resulted in the development of new targeted therapeutics which use revealed that the antiproliferative effects of COP1 blockade suppressed neoplastic growth in both wild-type and mutant p53-containing liver tumors, without unwanted immune responses.

2. AIMS

In human keratinocytes, the role of huCOP1 protein has not yet been characterized, therefore we aimed

- to characterize huCOP1 expression in cultured human keratinocytes and in human epidermis,
- to investigate the effect of UVB on huCOP1 expression in human cultured keratinocytes and human epidermis,
- to elucidate the COP1-p53 interaction in human keratinocytes,
- to determine the effect of UVB in COP1-silenced human keratinocytes
- to characterize changes in COP1 expression during keratinocyte differentiation.

3. MATERIALS AND METHODS

3.1. Human tissue samples. Culturing of normal human keratinocytes and HaCaT keratinocytes

Human tissue samples from various human organs were taken from patients who underwent different operations at the Department of Surgery, University of Szeged. Only non-involved, healthy tissues were used for RNA isolation. All tissue samples were taken with the patients' informed consent and the approval of the local committee

Skin biopsies were obtained from healthy individuals undergoing plastic surgery; institutional approval and patient consent was given for experiments in adherence to the Helsinki guidelines. After removal of the subcutaneous tissue, skin biopsies were incubated overnight at 4 °C in Dispase solution (Roche). The epidermis was separated from the dermis,

and epidermal cells were prepared by using trypsin (0.25%). Human epidermal keratinocytes were seeded in serum-free Keratinocyte Basal Medium (Gibco) supplemented with L-glutamine and antibiotic/antimycotic solution containing penicillin, streptomycin and amphotericin B (Sigma). Keratinocytes were grown at 37 °C in a 5% CO₂ atmosphere.

For the differentiation of keratinocyte cultures, normal human keratinocytes were cultured until 90-95% confluence in the third passage. The culturing conditions were altered by increasing the calcium concentration in the culture media to 1.7 mM (in the form of CaCl₂). Samples were taken at 0, 1, 2, 4, 6, 8 and 10 days after switching to the high calcium concentration medium.

The spontaneously immortalized human keratinocyte cell line, HaCaT, kindly provided by Dr. N. E. Fusening (Heidelberg, Germany), was maintained in high-glucose DMEM (Gibco), with 10% fetal bovine serum (HyClone), supplemented with L-glutamine and antibiotic/antimycotic solution containing 100 U/l penicillin, 100 µl/ml streptomycin and amphotericin B (Sigma), at 37 °C in a 5% CO₂ atmosphere.

3.2. Immunocytochemistry, immunohistochemistry and immunofluorescence staining

Immunocytochemistry: Keratinocytes were grown on culture slides (BD Falcon). The slides were fixed and incubated with the primary rabbit anti-COP1 antibody (Bethyl), at a dilution of 1:5000; control slides were incubated with a mixture of COP1 Blocking peptide (Bethyl) and rabbit anti-COP1 antibody in a ratio of 3:1. After a rinse in TBS-Triton-X, sections were incubated with Alexa Fluor 488 labelled secondary anti-rabbit antibody (Invitrogen) (dilution 1:800) for 3 hs, and were then stained with DAPI to detect the nuclei. For detection and analysis, Tissue Facs (Tissue Gnostics) and an FV 1000 confocal microscope (Olympus) were used.

Immunohistochemical staining: Formalin-fixed, paraffin-embedded skin tissue sections were dewaxed. Nonspecific staining was prevented by preincubation, and the slides were then incubated overnight at 4 °C in a humid chamber with the primary antibody; rabbit anti-COP1 antibody was applied at 1:5000 dilution, and rabbit IgG (NeoMarkers) was used for isotype control staining. Slides were then incubated with a biotinylated secondary antibody (anti-rabbit IgG) at 1:800 dilution for 1 h at room temperature, followed by incubation with horse radish peroxidase-conjugated streptavidin for 1 h at room temperature (both from Vectastain ABC Kit), and at the end of the staining procedure peroxidase activity was detected by using 3,3-amino-9 ethylcarbazole (Sigma) as substrate. Slides were counterstained with hematoxylin (Sigma). Tissue staining was visualized with a Zeiss Axio Imager microscope and photographed with a PixelINK digital camera.

Immunofluorescence staining: was performed as described above in connection with the immunohistochemical staining until the incubation with the primary antibody. The slides were then incubated at 4 °C overnight in a humid chamber with the primary antibody; rabbit anti-COP1 antibody was applied at 1:4000 dilution; control slides were incubated with a mixture of COP1 Blocking peptide and rabbit anti-COP1 antibody in a ratio of 3:1. After that, sections were incubated with Alexa Fluor 488-labeled secondary anti-rabbit antibody (dilution 1:500) for 3 hs, and were then stained with DAPI to detect the nuclei. For detection and analysis, Tissue Facs (Tissue Gnostics) were used

3.3. Constructs

The pSUPER vector system (Oligoengine) was used for the transient expression of short interfering RNAs (siRNAs). Fifty-three-nt-long oligonucleotides of COP1 5' AGCTTcttgattggccaatgtcaTTCAAGAGAtgacattggccaaatcaagttC3' (sense) and COP1 5' AgaactaaaccggttacagtAAGTTCTCTactgtaaccggttagttcaaGAGCT3' (antisense) were cloned into the pSuperior.puro vector downstream of the H1 promoter.

3.4. Nucleofection

For silencing the expression of the COP1 gene, $7-8 \times 10^5$ cells were transiently transfected with a COP1 siRNA producing construct (siCOP1), and another group of cells received the empty pSUPER vector (Sp). DNA for transfection was purified with the Qiagen Plasmid Mini Kit (Qiagen). Transfection was carried out by nucleofection (Amaxa), and the Human Keratinocyte Kit (VPD-1002; Amaxa) was used for the nucleofection of keratinocytes. The transfection efficacy was determined by analyzing the expression of green fluorescent protein (GFP) with a FACScalibur flow cytometer and CellQuest software (Becton Dickinson) and the mean fluorescence intensity of the expressing construct was measured. The average efficacy of transfection was ~60 %.

3.5. UVB irradiation (FS20 lamp, excimer laser)

An FS20 lamp (Westinghouse) was used as a radiation source for UVB. To determine the highest nonlethal dose of UVB, we irradiated keratinocytes with 0, 10, 20, 40 and 60 mJ/cm² UVB light, and the viability of the treated cells was analyzed by MTT assay 24 and 48 hs after irradiation. The 40 mJ/cm² found to be the highest nonlethal UVB dose was therefore used in subsequent experiments. Samples for mRNA expression experiments were collected 12 hs after irradiation, while samples for protein expression experiments were collected 24 hs after irradiation.

The 308 nm excimer laser was utilized for human skin irradiation. The minimal erythema dose (MED) was determined with increasing doses of 308 nm UVB, and 2 MED was used for irradiation on a non-sun-exposed area of the body in subsequent experiments. Skin biopsies (6 mm) were taken from irradiated and non-irradiated skin 24 and 48 hs after irradiation with the patients' informed consent.

3.6. Reverse transcription and real-time RT-PCR

Total RNA was isolated from cell cultures with the TRIzol reagent (Life Technologies) 12 h after UVB irradiation, following the instructions of the manual. cDNA was generated with oligo(dT) and random hexamer primers from 1 µg of RNA, using the iScriptTM cDNA Synthesis Kit (Bio-Rad Laboratories) in a final volume of 20 µl. After reverse transcription, the real-time RT-PCR was performed to compare the abundance of COP1 mRNA using TaqMan[®] Gene Expression Assays (Applied Biosystems) for the RFD2 (cat. No. Hs00375437_m1) and for p53 (cat. No. Hs01034249_m1). The abundance of the gene of interest transcript was normalized to the expression of 18S ribosomal RNA for each examined sample. The primers for 18S were: 18S RNA forward: CCGCTACCACATCCAAGGAA, 18S RNA reverse: GCTGGAATTACCGCGGCT, 18S RNA TaqMan probe: TexRed-TGCTGGACCAGACTTGCCCTC-BHQ-1 (Integrated DNA Technologies, Coralville, IA, USA). The RT-PCR reactions were performed by using iQ Supermix in an iCycler (Bio-Rad).

3.7. Western blot analysis

Cells were washed twice with PBS, and protein was extracted by resuspending the cells in a solution containing 12.5% 0.5 mM Tris (pH 6.8), 2.5% 200 mM EDTA, 15% 10% SDS, 70% H₂O, 10 µl of protease inhibitor and 50 µl of β-mercaptoethanol per ml (all chemicals were obtained from Sigma-Aldrich) 24 h after UVB irradiation. Protein lysates were incubated for 5 min on ice and, after a 30-sec hard vortexing, cell debris was removed by centrifugation at 10 000 g for 10 min at 4 °C. For western-blot analysis, equal amounts of proteins were run on SDS-PAGE and then transferred to nitrocellulose membrane (Bio-Rad).

Membranes were blocked by incubation in Tris-buffered saline (150 mM NaCl, 25 mM Tris, pH 7.4) containing 0.05% Tween 20 (Sigma-Aldrich) and 3% non-fat dry milk (Fluka Chemie AG) for 2 hs at room temperature and subsequently incubated overnight at 4 °C with purified mouse monoclonal anti-human p53 antibody (Calbiochem), rabbit polyclonal anti-human COP1 antibody (Bethyl), and rabbit monoclonal anti-actin antibody (Sigma-Aldrich). Alkaline phosphatase-conjugated goat anti-mouse IgG (Sigma-Aldrich) and anti-rabbit IgG (Sigma-Aldrich) were used as secondary antibodies at 1:2000 dilution in the blocking buffer and membranes were incubated for 2 hs at room temperature. The blots were developed by using 5-bromo-4-chloro-3-indolyl phosphate/nitroblue tetrazolium as substrate (BCIP/NBT, Sigma-Aldrich).

4. RESULTS

4.1. COP1 expression and subcellular localization in cultured normal human keratinocytes and HaCaT cells

Only limited information is available on COP1 expression in normal human tissue samples, and we therefore set out to determine COP1 expression in a set of RNA samples isolated from various human organs. We were especially interested in whether COP1 was expressed in the epidermis. COP1 expression was detected in all of the studied organ types. Interestingly, the level of expression differed to a great extent in the different organs. The lowest level of COP1 mRNA was detected in the breast sample, in agreement with earlier literature data. The highest level was seen in the gallbladder, 16-fold higher than in the breast.

We also demonstrated by RT-PCR and western blot hybridization assays that normal human cultured keratinocytes express COP1 mRNA and protein. COP1 mRNA was detected both in normal human keratinocytes and in HaCaT keratinocytes. In HaCaT keratinocytes the amount of COP1 mRNA was higher. As demonstrated by western blot analysis, HaCaT cells also expressed higher levels of COP1 protein compared to normal human keratinocytes. To examine the role of COP1 as a regulator of p53 in keratinocytes, we used normal keratinocytes for our subsequent investigations. Since HaCaT cells express an extremely high and stable level of p53 protein, HaCaT cells did not seem to be a good model cell to study COP1-p53 interactions.

We next determined the subcellular localization of COP1 protein in cultured normal human keratinocytes. COP1 protein was detected both in the cytoplasm as well as in the nucleus in unstressed keratinocytes compared to control cells incubating with blocking peptid before staining. The bulk of the immunofluorescence staining of COP1 was restricted to the nucleus, and only mild positivity was visible in the cytoplasm.

4.2. COP1 and p53 expression in UVB-irradiated keratinocytes

COP1 mRNA expression decreases after UVB irradiation in keratinocytes. COP1 is a central negative regulator of photomorphogenesis in plants. In human keratinocytes permanently exposed to UVB, the role of huCOP1 has not yet been characterized. For further investigation of huCOP1 in epidermal cells, we performed *in vitro* and *in vivo* experiments. To determine the highest nonlethal dose of UVB, we irradiated keratinocytes with 0, 10, 20, 40 and 60 mJ/cm² UVB light. The dose-dependent changes in COP1 mRNA expression were detected and the viability of the irradiated cells was analyzed by MTT assay. The 40 mJ/cm² found to be the highest nonlethal UVB dose, 70-80% of the treated cells in the culture survived it, therefore we applied this dose for the irradiation of subconfluent keratinocyte cultures.

To measure the kinetic changes in COP1 mRNA expression, keratinocytes were harvested at various time points (0, 6, 12 and 24 hs) after exposure to 40 mJ/cm² UVB (n=3) and the COP1 mRNA expression was determined by real-time RT-PCR. The COP1 mRNA levels started to decline within 6 h after UVB exposure, were at an almost undetectable level after 12 hs, and then slowly increased, approaching the normal value at 24 h.

UVB irradiation decreases the COP1 protein level in keratinocytes. We determined the abundance of COP1 and p53 proteins in keratinocytes (n=6) at various time-points (0, 6, 12 and 24 hs) after irradiation with 40 mJ/cm² UVB. The UVB-induced a dose-dependent decreases in COP1 protein are shown in Fig. 10. In contrast with COP1 mRNA, COP1 protein reached its minimum level at 24 h after UVB treatment. The p53 protein level was already increased 6 hs after UVB exposure and remained at high levels throughout the 24-h experimental period.

To measure the kinetic changes in p53 mRNA expression, keratinocytes were harvested at various time points (0, 6, 12 and 24 hs) after exposure to 40 mJ/cm² UVB (n=3) and the p53 mRNA expression was determined by real-time RT-PCR. In contrast with the COP1 mRNA, the p53 mRNA levels were not significantly changed during 12 hs after irradiation, but a marked increase was apparent at 24 h after UVB exposure. These results were in agreement with previously published data on the UV-induced p53 expression in keratinocytes.

UVB irradiation alters the subcellular localization of COP1 protein in keratinocytes. Immunofluorescence staining of COP1 demonstrated that treatment with 40 mJ/cm² UVB affects the subcellular distribution of the protein. Following UVB treatment, a substantial proportion of the COP1 protein was detected in the cytoplasm. This was in stark contrast with the untreated cells, in which COP1 was predominantly localized in the nucleus. However, in the cytoplasm marked perinuclear COP1 staining in the UVB-irradiated keratinocytes was apparent. Semiquantitative analysis of immunofluorescence staining demonstrated that treatment with 40 mJ/cm² UVB affected the COP1 expression both in the nuclei and in the cytoplasm, to approximately the same extent.

4.3. Up-regulated p53 protein expression is further increased by UVB irradiation in gene-specifically silenced COP1 cells

Early data on COP1 expression in mammalian cells indicated that it is a critical negative regulator of p53 protein. UVB irradiation and COP1 silencing have an additive effect on the induction of p53 expression. To gain further insight into the p53 regulatory function of COP1 in keratinocytes, we determined p53 protein levels in cells in which the expression of COP1 was specifically silenced by shCOP1 RNA. Western-blot experiments revealed that the expression of shCOP1 RNA resulted in the depletion of COP1 protein and significant up-regulation of p53 protein as compared with the mock-transfected cells.

We next investigated the effect of UVB irradiation in shCOP1-transfected keratinocytes treated with 40 mJ/cm². In these experiments, we monitored the intracellular levels of COP1 and p53 proteins 24 hs after irradiation. We demonstrated that UVB irradiation further decreased the level of COP1 protein, whereas the expression of p53 exhibited an additional increase. We concluded that COP1 silencing and UVB irradiation have additive effects on p53 protein accumulation in normal human keratinocytes.

4.4. COP1 expression in normal and UVB-irradiated human skin

COP1 protein expression was detected in normal human skin with immunohistochemistry. COP1 expression was the most prevalent in the stratum granulosum of the epidermis, and mild positivity was also apparent in the basal layer. The staining was strongest in the cytoplasm of the epidermal keratinocytes.

The expression of COP1 protein in normal human skin was also investigated by immunofluorescence labeling. In agreement with the immunohistochemistry, we found mild cytoplasmic staining in the basal layer of the epidermis and strong cytoplasmic staining was present in the granular layer.

Next our aim was to study the effect of UVB irradiation on COP1 expression in human skin. Normal human skin was irradiated with 2 MED and skin biopsies were taken 24 and 48 hs after irradiation. Both immunohistochemistry and immunofluorescence stainings indicated changes in COP1 expression in UVB-irradiated normal human skin. The COP1 expression was already decreased in both the stratum basale and the stratum granulosum 24 hs after the irradiation and was even lower 48 hs following the irradiation. These results suggest that the COP1 expression changes are part of the long-term stress response of human skin in response to UVB irradiation.

4.5. The level of COP1 protein in differentiating keratinocytes

The pattern of COP1 expression in the human epidermis prompted us to further analyze the changes in expression of COP1 in *in vitro* differentiating keratinocytes and to investigate the COP1-p53 relationship during differentiation. For the modeling of keratinocyte differentiation, we cultured keratinocytes in serum-free keratinocyte medium, with the addition of 1.7 mM calcium. To monitor the differentiation of keratinocytes, the expression of involucrin, a well-known differentiation marker, was investigated through western blotting, and we found an extremely high overexpression of the protein in parallel with the aging of the culture. p53 protein was expressed at a low level in the differentiating keratinocytes and did not exhibit change with time in the experimental period. In contrast, the initiation of differentiation resulted in a definitive decrease in COP1 protein in the keratinocyte culture; this decrease was already obvious 2 days after the induction of the differentiation. In parallel with this, we detected similar changes in COP1 expression in cultured normal human keratinocytes without calcium addition: the COP1 expression displayed a gradual decrease in cultured keratinocytes.

5. DISCUSSION

COP1, the Constitutive Photomorphogenic Protein 1, is well conserved across species: it is essential for normal photomorphogenesis in plants, and it has also been detected in mammalian cells. COP1 acts as a multifunctional ubiquitin ligase in plants and targets key transcription factors for proteasome-dependent degradation. Oravec et al. reported that, in stark contrast to its negative regulatory function in the visible part of the spectrum, AtCOP1 acts as a positive regulator of UVB-induced signaling in plant cells, and UVB irradiation promotes the accumulation of AtCOP1 in the nucleus. These data indicate that AtCOP1 is a multifunctional E3 ligase, and both its subcellular distribution and its substrate specificity are wavelength-specific. The striking degree of structural conservation of huCOP1 with its *Arabidopsis* counterpart suggested the possibility of a functional similarity too. Similarly to AtCOP1, MmCOP1 and huCOP1 are also present in the nucleus and the cytoplasm of cells.

The initial characterization showed that MmCOP1 (rodent), like its plant counterpart, is indeed involved in ubiquitination and is itself a substrate of its own ubiquitination activity. The huCOP1 ubiquitin ligase has been found to promote ubiquitin-dependent p53 degradation directly by the 26S proteasome. This process is independent of Mdm2 or Pirh2, which are also known to interact with and negatively regulate p53. The expression of huCOP1 has been detected in many cell lines, but its role in human cells is still not well characterized.

Our aim was therefore to investigate in detail the expression and function of huCOP1 in keratinocytes, the cell type most exposed to UVB irradiation. Our investigations showed that COP1 protein is present in human keratinocytes and immortalized HaCaT keratinocytes. COP1 protein is predominantly expressed in the nucleus of keratinocytes under unstressed conditions, and milder positivity is detected in the cytoplasm. It was previously demonstrated by others that, in an unstressed condition, p53 also shows a predominantly nuclear localization, and it is monoubiquitinated by Mdm2. Following monoubiquitination, p53 may remain in the nucleus to be polyubiquitinated by Mdm2, Pirh2, COP1 or a complex of these proteins, or it may be exported to the cytoplasm. All these processes explain the high nuclear expression of COP1 protein. The nuclear export of p53 promotes its cytosolic accumulation where it is further polyubiquitinated by Mdm2, Pirh2, COP1 or a complex of these proteins and degraded in the cytosolic 26S proteasome.

The comparison of the COP1 mRNA levels in various human tissues revealed that the epidermis expressed a significantly lower level of COP1 mRNA than the colon or gallbladder, but it was higher as compared with the breast tissue, where it was almost undetectable. A weak COP1 signal in normal breast tissue has already been described, whereas in breast adenocarcinoma the overexpression of the protein was detected. The same was demonstrated in ovarian adenocarcinomas, where the positive COP1 signal was present only within the malignant cells, and the stromal compartment or normal ovarian tissues were negative. This is concomitant mainly with a decrease in steady-state wild-type p53 protein levels or p53-dependent transcription, indicating that one of the major roles of COP1 is to repress p53-dependent tumor suppression. The fact that we detected COP1 in the epidermis affirmed our aim of investigating its physiological functions in normal skin.

In order to elucidate the biological function of COP1 in the major constituent cell type of the epidermis, the keratinocytes, we investigated its expression in response to genotoxic stress such as UVB light. Although UVB light is the most prominent genotoxic stress for keratinocytes, its effect on COP1 expression has not yet been investigated in these cells. Previously, it was shown that COP1 protein expression is decreased following IR treatment in wild-type p53-expressing U2OS osteosarcoma cells and fibroblasts. It has also been demonstrated that a decreased abundance of COP1 protein can not be attributed to a decrease in COP1 mRNA level. Another previous investigation has shown that treatment with UV irradiation results in a COP1 protein decrease in NIH3T3 mouse fibroblast cells. In the present study, we found that the COP1 mRNA level exhibited a biphasic response to UVB. It was decreased early after UVB irradiation, and a slow increase was observed after 12 hs. In contrast with the biphasic mRNA profile, the COP1 protein level displayed a steady slow, but marked decrease. In a previous study, UVC stress induced an increase in COP1 protein level in both HeLa and U2OS cells, although the COP1 mRNA showed a biphasic response characterized by a downregulation at early timepoints followed by a subsequent upregulation. These observations indicate that UV irradiation differentially affects COP1 protein levels in human keratinocytes and cancer cells.

Several data suggest that, besides its expression changes, the function of COP1 may also be altered by its intracellular movement. The first data on COP1 intracellular localization revealed that UVB induces the subcellular redistribution of AtCOP1. However, in *Arabidopsis thaliana*, UVB increased the accumulation of AtCOP1 in the nuclei and AtCOP1 has been postulated to regulate UVB responses positively. This is in marked contrast with mammalian cells, in which the immunofluorescence staining of COP1 indicated intensive positivity, mainly in the nucleus, in the absence of UVB. A similar subcellular redistribution of COP1 protein was also detected: ionizing radiation results in the nuclear export of COP1 protein in Flag-COP1-transfected H1299 cells, the primarily nuclear COP1 staining becoming cytosolic after IR. In line with the above findings, our results showed that UVB irradiation resulted in a similar decrease in COP1 protein both in the nuclei and in the cytoplasm, and a marked amount of the protein was centered in the perinuclear region of the cytoplasm. This means that in normal keratinocytes UVB irradiation caused not only a significant decrease in the COP1 protein level, but also a notable subcellular redistribution.

COP1 is a negative regulator of p53, and the depletion of COP1 by siRNA therefore stabilizes p53: COP1 silencing resulted in a higher p53 protein level in unstressed U2OS cells. In agreement with this, in our experimental set-up we saw that the depletion of COP1 by siRNA in normal keratinocytes resulted in an increase in p53 steady-state levels in unstressed conditions, indicating that COP1 has a negative regulatory role on p53 in keratinocytes too. We investigated the COP1-p53 connection in UVB-induced keratinocytes and found that the UVB irradiation of COP1-silenced keratinocytes resulted in an enhanced accumulation of p53 protein that was evident 24 h after the irradiation. COP1 silencing and UVB irradiation exerted an additive effect on p53 protein accumulation. This indicates that depletion of COP1 stabilizes p53 protein in both unstressed cells and UVB-irradiated cells and that the lack of COP1 may sensitize keratinocytes to UVB-induced cell death. Taken together, these results suggest that COP1 has an important role in the UVB response and fine tuning of apoptotic processes of keratinocytes by regulating p53.

We saw that huCOP1 contributes to the UV-induced stress response of keratinocytes. Next, we set out to investigate whether this molecule has any role in the regulation of keratinocyte differentiation and proliferation. We found that COP1 is highly expressed in normal human skin, in both the basal and the granular layers. This suggests that COP1 may have a role in the early steps of differentiation, in which the decision occurs as to whether the basal proliferating cell will enter the proliferation program. HuCOP1 is also expressed in the granular layer of the epidermis, in cells that are in the final stages of differentiation before cell death and cornification. The pleiotropic nature of huCOP1 and its role in complex regulatory functions could be revealed when huCOP1 and p53 expressions were studied in different layers of the normal epidermis upon UVB irradiation. p53 levels are low in the normal epidermis, and its induction results in the apoptosis of proliferating keratinocytes; as keratinocytes differentiate, they become insensitive to p53-induced apoptosis. It is well known that UVB light induces p53 in the proliferative compartment of the basal layer, resulting in protection from the tumorigenic effects of DNA damage; in our experiments we detected the opposite: decreases of the huCOP1 level in both the basal and the granular layer.

Consistent with the huCOP1 staining in the epidermis, differentiating keratinocytes express COP1 in the proliferative stage. As keratinocytes lose their proliferation potential and begin to express differentiation markers such as involucrin, K1 and K10, the COP1 expression undergoes a gradual decrease. This decrease was not associated with any changes in the level of p53; in fact, the p53 level did not show marked changes through differentiation.

These *in vitro* data are in agreement with the immunohistochemical data that huCOP1 is decreased in the suprabasal layer as compared with the basal layer of keratinocytes. The gradual decrease of COP1 protein in differentiating keratinocytes indicates that indeed COP1 is important through p53 regulation in the induction of keratinocytes apoptosis. The reason why fully differentiated, non-proliferative keratinocytes, which do not express p53, still contain COP1 is not clear. Further studies are needed to determine whether COP1 has other functions.

Our *in vitro* and *in vivo* data suggest that huCOP1 has a role in the fine tuning of UVB response in keratinocytes via regulation of the p53 function. The high level of huCOP1 in the basal layer of the epidermis, a UV-sensitive proliferating layer, is also an important observation in our experiments, suggesting that huCOP1 has an important role and may function as an oncogene which promotes tumorigenesis in p53-expressing cells, and that interventions targeting huCOP1 could well provide new opportunities in cancer therapy.

LIST OF PUBLICATIONS

I. Ágnes Kinyó, Zsuzsanna Kiss-László, Szabolcs Hambalkó, Attila Bebes, Mária Kiss, Márta Széll, Zsuzsanna Bata-Csörgő, Ferenc Nagy and Lajos Kemény: COP1 Contributes to UVB-Induced Signaling in Human Keratinocytes. *Journal of Investigative Dermatology* 130, 541–545; (2010) **IF: 5,543**

II. Kemény Lajos dr., Kinyó Ágnes dr., Hambalkó Szabolcs, Bebes Attila, Kiss Mária dr., Polyánka Hilda, Kiss-László Zsuzsanna dr., Bata-Csörgő Zsuzsanna dr., Nagy Ferenc dr., Széll Márta dr.: A COP1 az UVB-indukált jelátviteli út tagja humán keratinocitákban. *Bőrgyógyászati és Venereológiai Szemle* 85 (2), 49-54; (2009)

III. Kemény Lajos dr., Kinyó Ágnes dr., Baltás Eszter dr., Csoma Zsanett dr., Dósa Piroska dr., Gyulai Rolland dr., Bata Zsuzsanna dr., Dobozy Attila dr. A 308 nm-es excimer lézer a psoriasis kezelésében. *Bőrgyógyászati és Venereológiai Szemle* 82 (2), 67-71. (2006)

Publications indirectly related to the subject of the dissertation

1. Kinyó Ágnes dr., Gyulai Rolland dr., Varga Erika dr., Korom Irma dr., Bata-Csörgő Zsuzsanna dr., Kemény Lajos dr., Allopurinol-okozta hiperszenzitivitási szindróma. *Bőrgyógyászati és Venereológiai Szemle* 84 (3) 86-89. (2008)

2. Kinyó Ágnes, Belső Nóra, Nagy Nikoletta, Pálvölgyi Attila, Nagy István, Korom Irma, Varga Erika, Kemény Lajos, Bata-Csörgő Zsuzsanna. Strontium Ranelate-induced DRESS Syndrome with Persistent Autoimmune Hepatitis. *Acta Derm Venereol.* 2011 Jan 17. doi: 10.2340/00015555-1014. [Epub ahead of print] No abstract available.

IF: 3,007

7. ACKNOWLEDGMENTS

I would like to thank to Prof. Dr. Lajos Kemény for the possibility to perform these studies in the laboratory of the Department of Dermatology and Allergology, University of Szeged, and for his supervision of this work. Special thanks are due to Prof. Dr. Ferenc Nagy for his excellent suggestions and invaluable advice.

I thank Prof. Dr. Márta Széll for her constant support and advice, Dr. Zsuzsanna Kiss László and Dr. Mária Kiss for their support, and Prof. Dr. Zsuzsanna Bata-Csörgő for her valuable discussions.

I am grateful to my colleagues Szabolcs Hambalkó and Attila Bebes. I further thank Bernadett Kormos, Andrea Tanácsné Bajkán and Hilda Polyánka for their help in the preparative work. Special thanks are also due to Dr. Nikoletta Nagy for her constant support.

I am likewise very grateful to all colleagues who have provided me with so much help in my work.

This work was supported by the TÁMOP-4.2.2-08/1. and TÁMOP 4.2.1.B grants, and by the L'Oréal Inc (Hungary). The Project named „TÁMOP-4.2.1/B-09/1/KONV-2010-0005 – Creating the Center of Excellence at the University of Szeged” is supported by the European Union and co-financed by the European Regional Fund.



www.nfu.hu